



# Letters in Nonlinear Analysis and its Application

Peer Review Scientific Journal  
ISSN: 2958-874x

## Three-point fixed point results in $b$ -metric spaces

Cristina Maria Păcurar<sup>a,\*</sup>

<sup>a</sup>Faculty of Mathematics and Computer Science, Transilvania University of Braşov, 50 Iuliu Maniu Blvd., Braşov, Romania

---

### Abstract

In this paper, we explore fixed-point theorems in  $b$ -metric spaces, focusing on mappings that contract the perimeters of triangles and their generalizations. We establish new fixed-point results for such mappings under specific conditions, demonstrating their existence and uniqueness in  $b$ -metric spaces. Additionally, we introduce three-point Kannan-type generalized mappings, extending classical fixed-point theorems to a broader context. These results not only generalize existing theorems but also open new directions for further research in fixed-point theory within the  $b$ -metric framework.

*Keywords:* Fixed point,  $b$ -metric, mappings contracting perimeters of triangles  
*2010 MSC:* 47H10, 54E50, 47S20

---

### 1. Introduction

The concept of  $b$ -metric spaces was introduced by S. Czerwik (see [5]) in 1993 to generalize the traditional notion of metric spaces. The concept of  $b$ -metric spaces has since garnered significant attention in the field of mathematical analysis, particularly in the study of fixed-point theorems and their applications (see [4, 6, 7, 11, 12, 13] and the references cited herein). An interesting exposition of the early developments of the concept can be found in [2].

Very recently, Petrov introduced in [16] a new type of mappings called mappings contracting perimeters of triangles, which are a three-point analogue of Banach contractions [1]:

---

\*Corresponding author

Email address: [cristina.pacurar@unitbv.ro](mailto:cristina.pacurar@unitbv.ro) (Cristina Maria Păcurar )

**Definition 1.1** (Petrov [16]). Let  $(X, d)$  be a metric space with  $|X| \geq 3$ . We shall say that  $T: X \rightarrow X$  is a mapping contracting perimeters of triangles on  $X$  if there exists  $\alpha \in [0, 1)$  such that the inequality

$$d(Tx, Ty) + d(Ty, Tz) + d(Tz, Tx) \leq \alpha[d(x, y) + d(y, z) + d(z, x)],$$

holds for all three pairwise distinct points  $x, y, z \in X$ .

Petrov proved in [16] a fixed point theorem for these kind of mappings:

**Theorem 1.2** (Petrov [16]). *Let  $(X, d)$ ,  $|X| \geq 3$  be a complete metric space and let  $T: X \rightarrow X$  be a mapping contracting perimeters of triangles on  $X$ . Then,  $T$  has a fixed point if and only if  $T$  does not possess periodic points of prime period 2. The number of fixed points is at most 2.*

The newly introduced mappings were further studied and extended in [3, 8, 9, 14, 15, 17, 18, 19].

## 2. Preliminaries

Let us recall some basic notions related to  $b$ -metric spaces. We make the notation  $\mathbb{R}^+ = [0, \infty)$ .

Let  $d: X \times X \rightarrow \mathbb{R}^+$  and  $s \geq 1$ . We say that  $d$  is a  $b$ -metric on  $X$ , if for all  $x, y \in X$ , we have

(d<sub>1</sub>)  $d(x, y) = 0$  if and only if  $x = y$ ;

(d<sub>2</sub>)  $d(x, y) = d(y, x)$ ;

(d<sub>3</sub>)  $d(x, y) \leq s[d(x, z) + d(z, y)]$ .

**Definition 2.1.** Let  $(X, d)$  be a  $b$ -metric. A sequence  $(x_n)$  in  $X$  is said to be

- a.  *$b$ -convergent* to a point  $x \in X$  if for every  $\epsilon > 0$ , there exists a positive integer  $N$  such that for all  $n \geq N$ ,

$$d(x_n, x) < \epsilon.$$

- b.  *$b$ -Cauchy sequence* if for every  $\epsilon > 0$ , there exists a positive integer  $N$  such that for all  $m, n \geq N$ ,

$$d(x_n, x_m) < \epsilon.$$

**Definition 2.2.** Let  $(X, d)$  be a  $b$ -metric space. The space  $(X, d)$  is said to be  *$b$ -complete* if every  $b$ -Cauchy sequence in  $X$  is  $b$ -convergent in  $X$ .

In [13], the following lemma related to  $b$ -Cauchy sequences was proved:

**Lemma 2.3** ([13]). *Every sequence  $(x_n)_{n \in \mathbb{N}}$  of elements from a  $b$ -metric space  $(X, d)$  of constant  $s$ , having the property that there exists  $\gamma \in [0, 1)$  such that*

$$d(x_{n+1}, x_n) \leq \gamma d(x_n, x_{n-1}),$$

for every  $n \in \mathbb{N}$ , is  $b$ -Cauchy.

### 3. Mappings contracting perimeters of triangles in b-metric spaces

**Definition 3.1.** Let  $(X, d)$  be a b-metric space with  $|X| \geq 3$ . We shall say that  $T: X \rightarrow X$  is a mapping contracting perimeters of triangles in b-metric spaces if there exists  $\lambda \in [0, \frac{1}{s})$  such that the inequality

$$d(Tx, Ty) + d(Ty, Tz) + d(Tz, Tx) \leq \lambda[d(x, y) + d(y, z) + d(z, x)], \tag{1}$$

holds for all three pairwise distinct points  $x, y, z \in X$ .

**Theorem 3.2.** Let  $(X, d)$  be a complete b-metric space with  $|X| \geq 3$  and let the mapping  $T: X \rightarrow X$  satisfy the following two conditions:

- (i)  $T(Tx) \neq x$  for all  $x \in X$  such that  $Tx \neq x$ ;
- (ii)  $T$  is a mapping contracting perimeters of triangles in b-metric spaces.

Then,  $T$  has a fixed point. The number of fixed points is at most two.

*Proof.* Let  $x_0 \in X$ , arbitrarily chosen, but fixed and the Picard iteration

$$x_{n+1} = Tx_n, \quad \forall n \geq 0.$$

We shall show that  $T$  has at least one fixed point. Suppose that  $x_n$  is not a fixed point of the mapping  $T$  for every  $n = 0, 1, \dots$ . Then, we have  $x_n = Tx_{n-1} \neq x_{n-1}$  and  $x_{n+1} = T(Tx_{n-1}) \neq x_{n-1}$  for every  $n = 1, 2, \dots$ . Hence, by condition (i),  $x_{n-1}, x_n$  and  $x_{n+1}$  are pairwise distinct. Then taking in (1)  $x = x_{n-1}, y = x_n, z = x_{n+1}$  we obtain

$$\begin{aligned} d(Tx_{n-1}, Tx_n) + d(Tx_n, Tx_{n+1}) + d(Tx_{n+1}, Tx_{n-1}) &\leq \\ &\leq \lambda[d(x_{n-1}, x_n) + d(x_n, x_{n+1}) + d(x_{n+1}, x_{n-1})]. \end{aligned} \tag{2}$$

For every  $n = 0, 1, \dots$ , let

$$p_n = d(x_n, x_{n+1}) + d(x_n, x_{n+2}) + d(x_{n+1}, x_{n+2}).$$

Then, (2) becomes  $p_n \leq \lambda p_{n-1}$  for every  $n = 1, 2, \dots$ . Hence, by induction, we get

$$p_n \leq \lambda^n p_0. \tag{3}$$

Now, let  $p \in \mathbb{N}, p \geq 1$ . By (d<sub>3</sub>) we have

$$\begin{aligned} d(x_n, x_{n+p}) &\leq s[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+p})] \leq \\ &\leq sd(x_n, x_{n+1}) + s^2[d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_{n+p})] \leq \\ &\dots\dots\dots \\ &\leq sd(x_n, x_{n+1}) + s^2d(x_{n+1}, x_{n+2}) + \dots + s^p d(x_{n+p-1}, x_{n+p}) \leq \\ &\stackrel{(3)}{\leq} s\lambda^n p_0 + s^2\lambda^{n+1} p_0 + \dots + s^p \lambda^{n+p-1} p_0 = \\ &= p_0 s \lambda^n (1 + s\lambda + \dots + s^{p-1} \lambda^{p-1}) \leq \\ &\stackrel{s\lambda < 1}{\leq} p_0 s \lambda^n \frac{1}{1 - s\lambda}. \end{aligned}$$

Now, since  $s\lambda < 1$ , passing to limit as  $n \rightarrow \infty$ , we obtain that  $\lim_{n \rightarrow \infty} d(x_n, x_{n+p}) = 0$ , which implies that  $\{x_n\}$  is a **b**-Cauchy sequence. By  $b$ -completeness of  $(X, d)$ , we obtain that  $\{x_n\}$  is  $b$ -convergent and we get that  $\{x_n\}$  has a limit  $x^* \in X$ . Let us prove that  $Tx^* = x^*$ .

Since  $x_n, x_{n+1}$  and  $x_{n+2}$  are pairwise distinct for every  $n = 0, 1, \dots$ , there exists a subsequence  $\{x_{n(k)}\}_{k \geq 0}$  such that  $x_{n(k)}, x_{n(k)+1}$  and  $x^*$  are pairwise distinct for every  $k = 0, 1, \dots$ . Indeed, if  $x^*$  does not belong to the sequence  $\{x_{n(k)}\}$ , then since  $x_{n(k)} \neq Tx_{n(k)} = x_{n(k)+1}$ , the points  $x_{n(k)}, x_{n(k)+1}$  and  $x^*$  are pairwise distinct. If  $x^*$  belongs to the sequence  $\{x_{n(k)}\}$ , then since  $x_{n(k)} = Tx_{n(k)-1} \neq x_{n(k)-1}$  and  $x_{n(k)+1} = T(Tx_{n(k)}) \neq x_{n(k)}$  for every  $k = 0, 1, \dots$ , the points  $x_{n(k)-1}, x_{n(k)}, x_{n(k)+1}$  are pairwise distinct. Suppose  $k$  is the smallest possible index for which  $x^* = x_{n(k)-1}$  then  $x_{n(k)}, x_{n(k)+1}$  and  $x^*$  are pairwise distinct for every  $k = 0, 1, \dots$ .

Now, taking in (1)  $x = x_{n(k)}, y = x_{n(k)+1}$  and  $z = x^*$ , we obtain

$$d(Tx_{n(k)}, Tx_{n(k)+1}) + d(Tx_{n(k)+1}, Tx^*) + d(Tx^*, Tx_{n(k)}) \leq \lambda[d(x_{n(k)}, x_{n(k)+1}) + d(x_{n(k)+1}, x^*) + d(x^*, x_{n(k)})],$$

so ,

$$d(x_{n(k)+1}, x_{n(k)+2}) + d(x_{n(k)+2}, Tx^*) + d(Tx^*, x_{n(k)+1}) \leq \lambda[d(x_{n(k)}, x_{n(k)+1}) + d(x_{n(k)+1}, x^*) + d(x^*, x_{n(k)})].$$

Taking the limit as  $k \rightarrow \infty$  we get

$$2d(x^*, Tx^*) \leq 0,$$

by where  $d(x^*, Tx^*) = 0$ , so  $x^*$  is a fixed point of  $T$ .

Now, suppose that there exist three distinct fixed point of  $T, x^*, y^*, z^* \in X$ , then, by (1) we obtain

$$d(Tx^*, Ty^*) + d(Ty^*, Tz^*) + d(Tz^*, Tx^*) \leq \lambda[d(x^*, y^*) + d(y^*, z^*) + d(z^*, x^*)],$$

so

$$(1 - \lambda)[d(x^*, y^*) + d(y^*, z^*) + d(z^*, x^*)] \leq 0,$$

which implies  $1 - \lambda < 0$ , which is a contradiction, so  $T$  has at most two fixed points. □

**Example 3.3.** Let  $X = \{0, 1, 2\}$  and  $d(0, 0) = d(1, 1) = d(2, 2) = 0, d(1, 0) = d(0, 1) = \frac{1}{2}, d(2, 0) = d(0, 2) = 5$  and  $d(2, 1) = d(1, 2) = 6$ . Then,  $d$  is a  $b$ -metric, but it is not a metric as

$$d(2, 1) = 6 > 5.5 = d(2, 0) + d(0, 1).$$

Let  $T: X \rightarrow X$  such that  $T0 = T2 = 0$  and  $T1 = 1$ . Then  $T$  satisfies (1) for  $\lambda > \frac{2}{23}$  as

$$d(T0, T1) + d(T1, T2) + d(T2, T0) = 2d(0, 1) = 1 \leq 11.5 = d(0, 1) + d(1, 2) + d(2, 0).$$

Let us also note that  $T(Tx) \neq x$  for every  $x \neq Tx$ , and  $T$  has two fixed points.

#### 4. Three point Kannan generalized mappings in b-metric spaces

**Definition 4.1.** Let  $(X, d)$  be a  $b$ -metric space with  $|X| \geq 3$ . We shall say that  $T: X \rightarrow X$  is a three point Kannan generalized mapping in  $b$ -metric spaces if there exists  $\lambda \in [0, \frac{1}{2})$  such that the inequality

$$d(Tx, Ty) + d(Ty, Tz) + d(Tz, Tx) \leq \lambda[d(x, Tx) + d(y, Ty) + d(z, Tz)], \tag{4}$$

holds for all three pairwise distinct points  $x, y, z \in X$ .

**Theorem 4.2.** Let  $(X, d)$  be a complete  $b$ -metric space with  $|X| \geq 3$  and let the mapping  $T: X \rightarrow X$  satisfy the following two conditions:

- (i)  $T(Tx) \neq x$  for all  $x \in X$  such that  $Tx \neq x$ ;
- (ii)  $T$  is a three point Kannan generalized mapping in  $b$ -metric spaces.

Then,  $T$  has a fixed point. The number of fixed points is at most two.

*Proof.* Let  $x_0 \in X$ , arbitrarily chosen, but fixed and the Picard iteration

$$x_{n+1} = Tx_n, \quad \forall n \geq 0.$$

We shall show that  $T$  has at least one fixed point. As in the proof of theorem 3.2, we suppose that  $x_n$  is not a fixed point of the mapping  $T$  for every  $n = 0, 1, \dots$ . Then, we have  $x_n = Tx_{n-1} \neq x_{n-1}$  and  $x_{n+1} = T(Tx_{n-1}) \neq x_{n-1}$  for every  $n = 1, 2, \dots$  and by condition (i),  $x_{n-1}, x_n$  and  $x_{n+1}$  are pairwise distinct. Then taking in (4)  $x = x_{n-1}, y = x_n, z = x_{n+1}$  we obtain

$$\begin{aligned} d(Tx_{n-1}, Tx_n) + d(Tx_n, Tx_{n+1}) + d(Tx_{n+1}, Tx_{n-1}) &\leq \\ &\leq \lambda[d(x_{n-1}, Tx_{n-1}) + d(x_n, Tx_n) + d(x_{n+1}, Tx_{n+1})], \end{aligned} \tag{5}$$

so

$$\begin{aligned} d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + d(x_{n+2}, x_n) &\leq \\ &\leq \lambda[d(x_{n-1}, x_n) + d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2})], \end{aligned}$$

by which

$$(1 - \lambda)[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2})] + d(x_{n+2}, x_n) \leq \lambda d(x_{n-1}, x_n).$$

Since

$$(1 - \lambda)d(x_n, x_{n+1}) \leq (1 - \lambda)[d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2})] + d(x_{n+2}, x_n),$$

we obtain

$$(1 - \lambda)d(x_n, x_{n+1}) \leq \lambda d(x_{n-1}, x_n). \tag{6}$$

Since  $\lambda < \frac{1}{2}$ , we have  $\gamma = \frac{\lambda}{1 - \lambda} \in [0, 1)$ . Then, (6) becomes

$$d(x_n, x_{n+1}) \leq \gamma d(x_{n-1}, x_n),$$

so by Lemma 2.3, we obtain that  $\{x_n\}$  is a  $b$ -Cauchy sequence, and by completeness of  $X$ , we obtain that there exists  $x^* \in X$  such that  $\lim_{n \rightarrow \infty} x_n = x^*$ .

Now let us prove that  $Tx^* = x^*$ . As in the proof of theorem 3.2 there exists a subsequence  $\{x_{n(k)}\}_{k \geq 0}$  such that  $x_{n(k)}, x_{n(k)+1}$  and  $x^*$  are pairwise distinct for every  $k = 0, 1, \dots$

Now, taking in (4)  $x = x_{n(k)}, y = x_{n(k)+1}$  and  $z = x^*$ , we obtain

$$\begin{aligned} d(Tx_{n(k)}, Tx_{n(k)+1}) + d(Tx_{n(k)+1}, Tx^*) + d(Tx^*, Tx_{n(k)}) &\leq \\ &\leq \lambda[d(x_{n(k)}, Tx_{n(k)}) + d(x_{n(k)+1}, Tx_{n(k)+1}) + d(x^*, Tx^*)]. \end{aligned}$$

Hence,

$$\begin{aligned} d(x_{n(k)+1}, x_{n(k)+2}) + d(x_{n(k)+2}, Tx^*) + d(Tx^*, x_{n(k)+1}) &\leq \\ &\leq \lambda[d(x_{n(k)}, x_{n(k)+1}) + d(x_{n(k)+1}, x_{n(k)+2}) + d(x^*, Tx^*)]. \end{aligned}$$

Taking the limit as  $k \rightarrow \infty$  we get

$$2d(x^*, Tx^*) \leq \lambda d(x^*, Tx^*),$$

and since  $\lambda \in [0, \frac{1}{2})$  we obtain  $d(x^*, Tx^*) = 0$ , so  $x^*$  is a fixed point of  $T$ .

Now, suppose that there exist three distinct fixed point of  $T$ ,  $x^*, y^*, z^* \in X$ , then, by (4) we obtain

$$\begin{aligned} d(Tx^*, Ty^*) + d(Ty^*, Tz^*) + d(Tz^*, Tx^*) &\leq \\ &\leq \lambda[d(x^*, Tx^*) + d(y^*, Ty^*) + d(z^*, Tz^*)], \end{aligned}$$

so

$$d(x^*, y^*) + d(y^*, z^*) + d(z^*, x^*) \leq 0$$

which implies  $d(x^*, y^*) = d(y^*, z^*) = d(z^*, x^*) = 0$ , so  $T$  has at most two fixed points. □

**Example 4.3.** Let  $X = \{0, 1, 2\}$  and  $d(0, 0) = d(1, 1) = d(2, 2) = 0$ ,  $d(1, 0) = d(0, 1) = 10$ ,  $d(2, 0) = d(0, 2) = 2$  and  $d(2, 1) = d(1, 2) = 5$ . Then,  $d$  is a  $b$ -metric, but it is not a metric as

$$d(0, 1) = 10 > 7 = d(0, 2) + d(2, 1).$$

Let  $T: X \rightarrow X$  such that  $T0 = T1 = 0$  and  $T2 = 2$ . Then  $T$  satisfies (4) for  $\lambda > \frac{2}{5}$  as

$$d(T0, T1) + d(T1, T2) + d(T2, T0) = 2d(0, 2) = 4 \leq \lambda 10 = d(0, T0) + d(1, T1) + d(2, T2).$$

Let us also note that  $T(Tx) \neq x$  for every  $x \neq Tx$ , and  $T$  has two fixed points.

## References

- [1] S. Banach, *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fundamenta Mathematicae **3** (1922), 133-181.
- [2] V. Berinde, M. Pacurar, *The early developments in fixed point theory on  $b$ -metric spaces: a brief survey and some important related aspects*, Carpathian J. Math. **38** (2022), 523-538.
- [3] R. Bisht, E. Petrov, *A three point extension of Chatterjea's fixed point theorem with at most two fixed points*, arXiv:2403.07906.
- [4] M. Boriceanu, M. Bota, A. Petrusel, *Multivalued fractals in  $b$ -metric spaces*, Cent. Eur. J. Math. **8** (2010), 367-377.
- [5] S. Czerwik, *Contraction mappings in  $b$ -metric spaces*, Acta Math. Inform. Univ. Ostrav. **1** (1993), 5-11.
- [6] S. Czerwik, K. Dlutek, S. Singh, *Round-off stability of iteration procedures for set-valued operators in  $b$ -metric spaces*, J. Natur. Phys. Sci. **11** (2007), 87-94.
- [7] M. Jleli, B. Samet, C. Vetro, F. Vetro, *Fixed points for multivalued mappings in  $b$ -metric spaces*, Abstr. Appl. Anal. (2015), 718074.
- [8] M. Jleli, C.M. Pacurar, B. Samet, *New directions in fixed point theory in  $G$ -metric spaces and applications to mappings contracting perimeters of triangles*, arXiv preprint (2024) arXiv:2405.11648.
- [9] M. Jleli, C.M. Pacurar, B. Samet, *Fixed point results for contractions of polynomial type*, arXiv preprint (2024) arXiv:2406.03446
- [10] R. Kannan, *Some results on fixed points-II*, Amer. Math. Monthly. **76** (1969), 405-408.
- [11] E. Karapinar, *A short survey on the recent fixed point results on  $b$ -metric spaces*, Constr. Math. Anal. **1** (2018), 15-44.
- [12] W. Kirk, N. Shahzad,  *$b$ -Metric spaces. In: Fixed Point Theory in Distance Spaces*, pp. 113-131. Springer, Berlin (2014).
- [13] R. Miculescu, A. Mihail, *New fixed point theorems for set-valued contractions in  $b$ -metric spaces*, J. Fixed Point Theory Appl. **19** (2017), 2153-2163.
- [14] C. Pacurar, O. Popescu, *Fixed point theorem for generalized Chatterjea type mappings*, Acta Mathematica Hungarica **173** (2), 500-509.
- [15] C.M. Pacurar, O. Popescu, *Fixed points for three point generalized orbital triangular contractions*, arXiv preprint (2024) arXiv:2404.15682.
- [16] E. Petrov, *Fixed point theorem for mappings contracting perimeters of triangles*, J. Fixed Point Theory Appl. **25** (2023) 1-11.
- [17] E. Petrov, *Periodic points of mappings contracting total pairwise distance*, arXiv preprint (2024) arXiv:2402.02536.
- [18] E. Petrov, R. K. Bisht, *Fixed point theorem for generalized Kannan type mappings*, Rendiconti del Circolo Matematico di Palermo Series **2** (2024), 1-18.
- [19] O. Popescu, C.M. Pacurar, *Mappings contracting triangles*, arXiv preprint (2024) arXiv:2403.19488.